



Panther[®] SC

HERBICIDE



**LIQUID
FORMULA**

**FASTER TO MIX.
EASIER TO USE.**

Panther[®] SC: Pour on the Power

Panther[®] SC Herbicide provides excellent residual control of tough broadleaf weeds like marehail, Palmer amaranth, and waterhemp with speed and convenience. Offering flexible rotation options to many crops, Panther SC is a broad-spectrum pre-emerge broadleaf herbicide option for soybean, cotton and peanut growers. Plus, Panther SC fits into your production system - apply in the fall for a clean start in the spring, apply early preplant for burndown before planting, or apply preplant or pre-emerge for residual control in Roundup Ready[®] systems. Check the label for application timings and planting restrictions for various crops.

BROAD SPECTRUM WITH RESIDUAL CONTROL

- Controls a broad spectrum of annual broadleaf and grass weeds, including most broadleaf weeds resistant to glyphosate
- Convenient liquid formulation makes mixing and loading faster and easier

FLEXIBLE

- Offers long-lasting residual control to keep fields clean
- Flexible rotation options allow you to pick the crop that maximizes your profits
- Enhanced speed of control on a broader weed spectrum in labeled burndown programs

SAME RATE, SAME POWER

RATES ARE EQUIVALENT TO WDG FORMULATION:
1 oz Panther WDG = 1 fl oz Panther SC

EPA REG. NO. 71368-113

ACTIVE INGR. flumioxazin (44.0%)
(4 lb per gallon)

FORMULATION liquid

CHEM. FAMILY N-phenylphthalimide

GROUP NO. 14

MODE OF ACT. PPO inhibitor

REI 12 hours

SIGNAL WORD none

RES. USE no

PACKAGE SIZE 2.5 gal container (2 per case)



Nufarm

Grow a better tomorrow.



Panther SC

HERBICIDE

KEY USES

Cotton
 Dry beans
 Field corn
 Peanuts
 Soybeans
 Sugarcane
 Sweet potato
 Fallow land
 Bareground in non-crop farm areas

KEY WEEDS CONTROLLED

(See product label for complete list)

Carpetweed
 Chickweed (*common and mouseear*)
 Coffee senna
 Common ragweed
 Dandelion
 Eclipta
 Evening primrose, cutleaf
 False chamomile
 Florida beggarweed
 Florida pusley
 Golden crownbeard
 Hairy indigo
 Hemp sesbania
 Henbit
 Jimsonweed
 Kochia
 Lambsquarters, common
 Marestalk/horseweed
 Mustard (*tansy, tumble, wild*)
 Nightshades (*black, eastern black, hairy*)
 Palmer amaranth
 Pigweeds (*redroot, smooth, spiny amaranth, tumble*)
 Prickly sida (*teaweed*)
 Puncturevine
 Purslane (*common, horse*)
 Radish, wild
 Redmaids
 Redweed
 Shepherd's-purse
 Smallflower morningglory
 Spurred anoda
 Tropic croton
 Venice mallow
 Waterhemp (*common, tall*)
 Wild poinsettia

USE SUMMARY - ROTATION OPTIONS

RATE	CROP	ROTATIONAL INTERVAL
1 fl oz/A	Cotton (no-till or strip-till only)	14 days ¹
1.5 to 2 fl oz/A	Cotton (no-till or strip-till only)	21 days ¹
2 fl oz/A or less	Peanut, Soybean, Sugarcane and Sweet Potato	Immediately
	Field Corn (minimum and no-till)	14 days
	Cotton and Field Corn (conventional tillage), Rice, Sorghum, Sunflower, Tobacco and Wheat	30 days ¹
	Barley, Dry and Snap Beans, Flax, Pea, Rye, Safflower and Sweet Corn	3 mths
Up to 3 fl oz/A	Alfalfa, Canola, Clover, Oats, Potato, Sugar Beet and all other crops not listed ²	4 mths (tilled prior to planting) 8 mths (no tillage)
	Lentil	6 mths
	Peanut, Soybean, Sugarcane and Sweet Potato	Immediately
Up to 4 fl oz/A	Field Corn (minimum and no-till)	14 days
	Field Corn (conventional tillage) and Sorghum	30 days ¹
	Cotton, Rice, Sunflower, Tobacco and Wheat	2 mths ¹
	Barley, Dry and Snap Beans, Flax, Pea, Rye, Safflower and Sweet Corn	4 mths
	Alfalfa, Clover, Oat, Sugar Beet	5 mths (tilled prior to planting) 10 mths (no tillage)
	Canola and all other crops not listed ²	6 mths (tilled prior to planting) 12 mths (no tillage)
6 to 12 fl oz/A	Lentil	7 mths
	Sugarcane	Immediately
	Cotton, Field Corn, Peanut, Rice, Sorghum, Soybean, Sunflower, Tobacco and Wheat	4 mths
	Alfalfa, Canola, Sugar Beet and all other crops not listed ²	6 mths (tilled prior to planting) 12 mths (no tillage)
6 to 12 fl oz/A	Cotton, Field Corn, Peanut, Rice, Sorghum, Soybean, Sunflower, Tobacco and Wheat	9 mths
	Alfalfa, Canola, Sugar Beet and all other crops not listed ²	12 mths (tilled prior to planting) 18 mths (no tillage)
	Trees can be transplanted 2 months after an application of this product ³	

¹ At least 1" of rainfall/irrigation must occur between application and planting or crop injury may occur.

² Successful soil bioassay must be performed prior to planting crops.

³ Transplanted apple, apricot, avocado, bushberries (including blueberry), cherry, fig, grape, grapefruit, lemon, nectarine, nut trees (including pistachio), olive, orange, peach, pear, plum (including dried plum), and tangerine can be planted 2 months after application of 2 to 12 fl oz/A of this product.

For residual weed control, apply 2 to 4 fl oz per acre by ground or air depending upon crop and application timing. Use 10 to 30 gals of spray solution per acre for ground application in conventional tillage. Refer to the Panther® SC label for specific application instructions for burndown applications. To ensure thorough coverage in burndown applications, use 15 to 60 gals spray solution per acre. Use 20 to 60 gals per acre if dense vegetation or heavy crop residue is present. For aerial application as part of a burndown weed control program, apply Panther in 7 to 10 gals of water per acre. When used aerially for pre-emergence weed control, apply Panther in 5 to 10 gals of water per acre.

If weeds have emerged when application is made, use Panther in combination with a labeled burndown herbicide. Post-emergence control requires the addition of approved COC, MSO or NIS adjuvants.

See label for complete directions for use.



 NUFARMPPOINT.COM

For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit www.nufarm.com/us to download a full label.
 ©2018 Nufarm. Important: Always read and follow label instructions. Panther® is a trademark of Nufarm. Roundup Ready® is a registered trademark of Monsanto Technology LLC. 15-AG-0091-E