
Envoy Plus®

HERBICIDE



Take Control Over Problem Grasses in Public and Industrial Areas.

Envoy Plus® Herbicide is a tool for industrial vegetation managers looking for post-emergence control of grasses such as Johnsongrass, Bermudagrass, crabgrass and quackgrass in public areas. Envoy Plus takes out the problem grass while leaving non-target plants and trees unaffected.

ENVOY PLUS DELIVERS:

- Excellent post-emergence weed control
- Quickly translocates to take out the entire grass weed
- Broad-spectrum control of the toughest grasses Johnsongrass, Bermudagrass, barnyardgrass, crabgrass, foxtails, brome, wild oats, ryegrass and quackgrass
- Low-use rates for cost effectiveness and convenience
- No soil carryover

EPA REG. NO.	59639-132
ACTIVE INGR.	clethodim (12.6%)
FORMULATION	liquid
CHEM. FAMILY	cyclohexanedione
HRAC NO.	1
SIGNAL WORD	anutin n
JIGINAL WORD	caution
RESTRICTED	no
0.0.0.2.00.0	000000







KEY USES

Rights-of-way including;

dividers, highways, medians, pipelines, public utility lines, pumping stations, railroads, roads, substations, transformer stations

Airports, commercial buildings, electric utilities, fence lines, manufacturing plants, parkways, post-harvest croplands, rail yards, storage yards

KEY WEEDS CONTROLLED

(See product label for complete list)

Annual and perennial grasses

USE SUMMARY

APPLICATION INFORMATION

Apply Envoy Plus® post-emergence herbicide to actively growing grasses. Applications made to grass plants stressed by insufficient moisture or cold temperatures, or to grass plants exceeding recommended growth stages may result in unsatisfactory control. Do not apply under these conditions. In arid regions where irrigation is used to supplement limited rainfall, Envoy Plus should be applied as soon as possible after irrigation (within 7 days). In arid regions, a second application of Envoy Plus will generally provide more effective control of perennial grass weeds than a single application. Make second application to actively growing grass 2 to 3 weeks after emergence of new growth.

- Use a minimum spray volume of 5 gal/A for aerial applications and 15 gal/A for ground applications to fallow land. Apply only to actively growing grasses when the first grass reaches the recommended weed height.
- Annual grasses that emerge after the Envoy Plus application will not be controlled, and a second application may be necessary.
- The control of perennial grasses may require more than one application in non-tilled areas.
- Do not apply to grasses that have tillered, formed seedheads or exceeded recommended growth stage.

ANNUAL GRASS RATE	PERENNIAL GRASS RATE	USE INSTRUCTIONS
9 - 16 fl oz/A	12 - 32 fl oz/A	Do not apply more than 32 oz/A per application.
		Do not apply more than 64 oz/A (0.5 lb ai/A) per season.
		For repeat applications, make on a minimum of a 14 day interval.

Adjuvant: COC (1%) or NIS (0.25%). Apply to actively growing grass weeds. Treated grass stops growing soon after application and dies within 1 - 2 weeks. *Ammonium sulfate recommended for some uses (see label)*.

USE RESTRICTIONS

DO NOT apply if rain is expected within 1 hour of application as control may be unsatisfactory.

DO NOT apply a post-emergence broadleaf herbicide within one day following application of Envoy Plus or reduced grass control may result.

DO NOT apply under conditions of stress. Applying Envoy Plus under conditions that do not promote active grass growth will reduce herbicide effectiveness. These conditions include drought, excessive water, extremes in temperature, low humidity and grasses either partially controlled or stunted from prior pesticide applications. Grasses under these kinds of stressful conditions will not absorb and translocate Envoy Plus effectively, and will be less susceptible to herbicide activity.

Application on Long Island, New York, is restricted to no more than 32 fl oz of Envoy Plus (0.25 lb ai) per acre per season.

See label for complete directions for use.



